

CONTENTS

Special issue: Mass Spectrometry and N.m.r. Spectroscopy

Foreword	vii
Controlled reduction of acarbose: conformational analysis of acarbose and the resulting saturated products K. Bock, M. Meldal (Valby, Denmark), and S. Refn (Lyngby, Denmark)	1
A procedure for the analysis by mass spectrometry of the structure of oligosaccharides from high-mannose glycoproteins A.-S. Angel, P. Lipniunas, K. Erlansson, and B. Nilsson (Lund, Sweden)	17
N.m.r. studies of the conformation of analogues of methyl β -lactoside in methyl sulfoxide- d_6 A. Rivera-Sagredo, J. Jiménez-Barbero, and M. Martín-Lomas (Madrid, Spain)	37
The resolution into molecular species on desorption of glycolipids from thin-layer chromatograms, using combined thin-layer chromatography and fast-atom-bombardment mass spectrometry K.-A. Karlsson, B. Lanne, W. Pimlott, and S. Teneberg (Göteborg, Sweden)	49
Characterisation by ^1H -n.m.r. spectroscopy of oligosaccharides, derived from arabinoxylans of white endosperm of wheat, that contain the elements $\rightarrow 4)[\alpha\text{-L-Araf-(1}\rightarrow 3)]\text{-}\beta\text{-D-Xylp-(1}\rightarrow \text{or } \rightarrow 4)[\alpha\text{-L-Araf-(1}\rightarrow 2)]\text{-}\beta\text{-D-Xylp-(1}\rightarrow$ R. A. Hoffman, B. R. Leeflang, M. M. J. de Barse, J. P. Kamerling, and J. F. G. Vliegthart (Utrecht, The Netherlands)	63
One-bond carbon–proton coupling constants: angular dependence in α -linked oligosaccharides I. Tvaroska (Bratislava, Czechoslovakia) and F. R. Taravel (Grenoble, France)	83
N.m.r. and conformational analysis of the capsular polysaccharide from <i>Streptococcus pneumoniae</i> type 4 C. Jones, F. Currie, and M. J. Forster (South Mimms, Gt. Britain)	95
Solution conformation and dynamics of the octadeoxynucleotide d(CACTAGTG) ₂ : a multinuclear n.m.r. relaxation study A. N. Lane (London, Gt. Britain)	123
N.m.r. and conformational analysis of some 2,3-disubstituted methyl α -L-rhamnopyranosides N. K. Kochetkov, G. M. Lipkind, A. S. Shashkov, and N. E. Nifant'ev (Moscow, U.S.S.R.)	145
Characterisation of the N-linked oligosaccharides of the light chain of human glycoprotein IIb by f.a.b.-m.s. A. J. Reason, A. Dell, H. R. Morris (London, Gt. Britain), M. E. Rogers (West Chester, PA, U.S.A.), J. J. Calvete, and J. González-Rodríguez (Madrid, Spain)	169
Analysis of sialic acid-containing mucin oligosaccharides from porcine small intestine by high-temperature gas chromatography–mass spectrometry of their dimethylamides G. C. Hansson, J.-F. Bouhours, H. Karlsson (Göteborg, Sweden), and I. Carlstedt (Lund, Sweden)	179

Characterisation of minor tetra- to hepta-saccharides O-linked to human meconium glycoproteins by t.l.c-m.s. microsequencing of neoglycolipid derivatives in conjunction with conventional m.s. and ^1H -n.m.r. spectroscopy A. M. Lawson, E. F. Hounsell, M. S. Stoll (Harrow, Gt. Britain), J. Feeney (London, Gt. Britain), W. Chai, J. R. Rosankiewicz, and T. Feizi (Harrow Gt. Britain)	191
Collisional-activation tandem mass spectrometry of sodium adduct ions of methylated oligosaccharides: sequence analysis and discrimination between α -NeuAc-(2 \rightarrow 3) and α -NeuAc-(2 \rightarrow 6) linkages J. Lemoine, G. Strecker, Y. Leroy, B. Fournet, and G. Ricart (Villeneuve d'Ascq, France)	209
Characterisation of polysaccharides by in-source pyrolysis positive- and negative-ion direct chemical ionisation-mass spectrometry J. A. Lomax, J. J. Boon (Amsterdam, The Netherlands), and R. A. Hoffmann (Utrecht, The Netherlands)	219
Fast-atom-bombardment chemistry of sulfatide (3-sulfogalactosylceramide) Y. Ohashi and Y. Nagai (Tokyo, Japan)	235
<i>Notes</i>	
Measurement of the $J_{\text{C-1,H-1}}$ coupling constants for the <i>Escherichia coli</i> O1A O-polysaccharide, a comparison of some n.m.r. experiments A. Helander and L. Kenne (Stockholm, Sweden)	245
Linkage analysis in disaccharides by electrospray mass spectrometry D. Garozzo, G. Impallomeni, E. Spina (Catania, Italy), B. N. Green and T. Hutton (Altrincham, Gt. Britain)	253
Complete assignment of the ^1H - and ^{13}C -n.m.r. spectra of the O-deacetylated glucuronoxylomannan from <i>Cryptococcus neoformans</i> serotype B M. A. Skelton, H. van Halbeek (Athens, GA, U.S.A.), and R. Cherniak (Atlanta, GA, U.S.A.)	259
<i>Author index</i>	c1
<i>Subject index</i>	c3

